Roman Road species list – Birds

Compiled by David J. Barden Covering up to the end of 2009 Last updated 9 December 2012

Turdus merula Sylvia atricapilla Larus ridibundus Parus caeruleus Fringilla montifringilla Pyrrhula pyrrhula Bullfir Buteo buteo Branta canadensis Corvus corone Fringilla coelebs Chaff Corvus corone Carrio Fringilla coelebs Chaff Corvus canus Comr Emberiza calandra Corn Cuculus canorus Cuculus canorus Curle Charadrius morinellus Charadrius morinellus Curle Charadrius regulus Fieldf Sylvia borin Regulus regulus Goldc Pluvialis apricaria Carduelis carduelis	kbird kcap k-headed Gull Tit nbling inch card ada Goose ion Crow ifinch chaff ared Dove mon Gull n Bunting coo ew erel den Warbler lcrest len Plover	Notes 86% (PL) 16% (PL) 19% (PL) 84% (PL) Big flock mixed with Chaffinch in 2003 (NL8), Between 10 and 35 seen on 3 occasions in 2005 and 2007 (IW) 49% (PL), 2 males seen in 2006 (IW) 24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	Ref. (2000-2009) PL, ISGJ, BBBW PL, BRC, BBBW PL PL, BRC, ISGJ, BBBW IW, BRC PL, IW, BBBW PL, IW, BRC, NL8 PL, IW, BRC, ISGJ, BBBW, NL8 PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC PL, IW, BRC
Sylvia atricapilla Larus ridibundus Parus caeruleus Pringilla montifringilla Pyrrhula pyrrhula Bullfir Buteo buteo Branta canadensis Corvus corone Fringilla coelebs Chaff Corvus corone Carric Fringilla coelebs Chaff Corpus collybita Chiffo Streptopelia decaocto Collat Larus canus Comr Emberiza calandra Corn Cuculus canorus Cucken Numenius arquata Charadrius morinellus Charadrius morinellus Comr Charadrius regulus Prunella modularis Turdus pilaris Fieldf Sylvia borin Garde Regulus regulus Goldc Pluvialis apricaria Golde Carduelis carduelis Larus marinus Great Gull Otis tarda Creat Parus major Dendrocopos major Great Carduelis chloris Tringa nebularia Greer Tringa nebularia Greer Tringa nebularia Greer Tringa nebularia Greer Ardea cinerea Grey	kcap k-headed Gull Tit hbling inch rard ada Goose ion Crow ifinch chaff ared Dove mon Gull a Bunting koo ew erel den Warbler lcrest len Plover lfinch at Black-backed	16% (PL) 19% (PL) 84% (PL) Big flock mixed with Chaffinch in 2003 (NL8), Between 10 and 35 seen on 3 occasions in 2005 and 2007 (IW) 49% (PL), 2 males seen in 2006 (IW) 24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC, BBBW PL PL, BRC, ISGJ, BBBW IW, BRC PL, IW, BBBW PL, IW, BRC, NL8 PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC PL, ISGJ PL, ISGJ PL, ISGJ PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC PL, IW, BRC, ISGJ, BBBW PL, BRC PL, IW, BRC PL, IW, BRC
Larus ridibundus Parus caeruleus Pringilla montifringilla Pyrrhula pyrrhula Bullfir Buteo buteo Buzza Branta canadensis Corvus corone Carrio Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Larus canus Comr Emberiza calandra Cuculus canorus Cuculus canorus Cuculus canorus Curle Charadrius morinellus Corne Charadrius morinellus Corne Cuculus canorus Curle Charadrius morinellus Corne Cuculus canorus Curle Charadrius morinellus Curle Charadrius morinellus Corne Cuculus canorus Cucle Numenius arquata Curle Charadrius morinellus Curle Charadrius morinellus Corduelis apricaria Golde Carduelis chloris Carduelis ch	k-headed Gull Tit Inbling Inch Inch Inch Inch Inch Inch Inch Inch	19% (PL) 84% (PL) Big flock mixed with Chaffinch in 2003 (NL8), Between 10 and 35 seen on 3 occasions in 2005 and 2007 (IW) 49% (PL), 2 males seen in 2006 (IW) 24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL PL, BRC, ISGJ, BBBW IW, BRC PL, IW, BBBW PL, IW, BRC, NL8 PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC PL, IW, BRC, ISGJ, BBBW PL, BRC PL, IW, BRC PL, IW, BRC PL, IW, BRC
Parus caeruleus Fringilla montifringilla Brami Pyrrhula pyrrhula Bullfir Buteo buteo Buzza Branta canadensis Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Larus canus Emberiza calandra Corn Cuculus canorus Curle Charadrius morinellus Charadrius morinellus Dotte Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Pluvialis apricaria Golde Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Great Vooc Picus viridis Greer Carduelis chloris Tringa nebularia Greer Grey	Tit Inbling Inch Inch Inch Inch Inch Inch Inch Inch	Business and seem of the seem in 2005 (NL8), Between 10 and 35 seem on 3 occasions in 2005 and 2007 (IW) 49% (PL), 2 males seen in 2006 (IW) 24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 8% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC, ISGJ, BBBW IW, BRC PL, IW, BBBW PL, IW, BRC, NL8 PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC PL, ISGJ PL, ISGJ PL, ISGJ PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, IW, BRC, ISGJ, BBBW PL, IW, BRC PL, IW, BRC PL, IW, BRC PL, IW, BRC
Fringilla montifringilla Pyrrhula pyrrhula Bullfin Buteo buteo Branta canadensis Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Larus canus Emberiza calandra Corn Cuculus canorus Cunella adecaocto Cuclus canorus Curle Charadrius morinellus Dotte Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Golde Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Tringa nebularia Greet Tringa nebularia Greet Ardea cinerea Grey	ada Goose inch card ada Goose ion Crow iffinch ichaff ared Dove imon Gull in Bunting koo ew erel hock lifare len Warbler lcrest len Plover lfinch at Black-backed	Big flock mixed with Chaffinch in 2003 (NL8), Between 10 and 35 seen on 3 occasions in 2005 and 2007 (IW) 49% (PL), 2 males seen in 2006 (IW) 24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	IW, BRC PL, IW, BBBW PL, IW, BRC, NL8 PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL, IW, BRC, ISGJ, BBBW PL, IW, BRC, ISGJ, BBBW PL, IW, BRC PL, IW, BRC PL, IW, BRC PL, IW, BRC
Pyrrhula pyrrhula Bullfir Buteo buteo Branta canadensis Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Larus canus Comr Emberiza calandra Corn Cuculus canorus Numenius arquata Charadrius morinellus Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Carduelis chloris Greet Tringa nebularia Greet Tringa nebularia Greet Tringa nebularia Greet Grey	inch card ada Goose ion Crow iffinch ichaff ared Dove imon Gull in Bunting koo ew erel hock lifare len Warbler lcrest len Plover lfinch at Black-backed	seen on 3 occasions in 2005 and 2007 (IW) 49% (PL), 2 males seen in 2006 (IW) 24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC, NL8 PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, IW, BRC PL, IW, BRC PL, IW, BRC
Branta canadensis Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Larus canus Emberiza calandra Corn Cuculus canorus Numenius arquata Charadrius morinellus Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Carduelis chloris Greet Carduelis chloris Greet Tringa nebularia Greet Tringa nebularia Greet Grey	ada Goose ion Crow ifinch ichaff ared Dove imon Gull is Bunting koo ew erel inock iffare iden Warbler icrest ien Plover iffinch	49% (PL), 2 males seen in 2006 (IW) 24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC, NL8 PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, IW, BRC PL, IW, BRC PL, IW, BRC
Branta canadensis Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Larus canus Emberiza calandra Corn Cuculus canorus Numenius arquata Charadrius morinellus Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Carduelis chloris Greet Carduelis chloris Greet Tringa nebularia Greet Tringa nebularia Greet Grey	ada Goose ion Crow ifinch ichaff ared Dove mon Gull i Bunting koo ew erel hock lfare len Warbler lcrest len Plover lfinch at Black-backed	24%, seen in 1996, 1997, and frequently from 2003 onwards (PL), 1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW)	PL, IW, BRC, NL8 PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, IW, BRC PL, IW, BRC PL, IW, BRC
Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Collai Larus canus Comr Emberiza calandra Corn Cuculus canorus Cundus canorus Curle Charadrius morinellus Charadrius morinellus Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Goldc Carduelis carduelis Larus marinus Great Carduelis carduelis Goldf Carduelis carduelis Carduelis carduelis Goldf Carduelis carduelis Great Gull Otis tarda Parus major Carduelis chloris Great Tringa nebularia Greer Ardea cinerea Grey	ion Crow iffinch ichaff ared Dove imon Gull in Bunting ichaff ared Dove imon Gull in Bunting ichaff ared Dove imon Gull in Bunting ichaff ared ichaff	1 seen in 2005 (disturbed from feeding on rabbit) (IW); 2 seen in 2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW)	PL PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC
Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Collai Larus canus Comr Emberiza calandra Corn Cuculus canorus Cundus canorus Curle Charadrius morinellus Charadrius morinellus Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Goldc Carduelis carduelis Larus marinus Great Carduelis carduelis Goldf Carduelis carduelis Carduelis carduelis Goldf Carduelis carduelis Great Gull Otis tarda Parus major Carduelis chloris Great Tringa nebularia Greer Ardea cinerea Grey	ion Crow iffinch ichaff ared Dove imon Gull in Bunting ichaff ared Dove imon Gull in Bunting ichaff ared Dove imon Gull in Bunting ichaff ared ichaff	2006 (IW) 1%, only seen in June 2002 (PL) 36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW)	PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, IW, BRC
Corvus corone Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Collai Larus canus Comr Emberiza calandra Corn Cuculus canorus Cundus canorus Curle Charadrius morinellus Charadrius morinellus Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Regulus regulus Goldc Carduelis carduelis Carduelis carduelis Carduelis carduelis Gull Otis tarda Parus major Creat Vooc Picus viridis Great Carduelis chloris Greet Tringa nebularia Greet Ardea cinerea Grey	ion Crow iffinch ichaff ared Dove imon Gull in Bunting ichaff ared Dove imon Gull in Bunting ichaff ared Dove imon Gull in Bunting ichaff ared ichaff	36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW)	PL, ISGJ PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, IW, BRC
Fringilla coelebs Chaff Phylloscopus collybita Streptopelia decaocto Collar Larus canus Comr Emberiza calandra Cuculus canorus Numenius arquata Charadrius morinellus Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Carduelis carduelis Goldet Larus marinus Parus major Dendrocopos major Great Wooc Picus viridis Greer Carduelis chloris Tringa nebularia Greer Ardea cinerea Grey	iffinch ichaff ared Dove imon Gull is Bunting icoo ew erel inock Iffare Iden Warbler Icrest Iden Plover Iffinch at Black-backed	36% (PL) 94% (PL); big flock mixed with Brambling in 2003 (NL8); part of a mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW)	PL, BRC, ISGJ, BBBW, NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, BRC
Phylloscopus collybita Chiffo Streptopelia decaocto Collar Larus canus Comr Emberiza calandra Corn Cuculus canorus Cucke Numenius arquata Curle Charadrius morinellus Dotte Prunella modularis Dunn Turdus pilaris Fieldf Sylvia borin Garde Regulus regulus Goldo Pluvialis apricaria Golde Carduelis carduelis Goldf Larus marinus Great Gull Otis tarda Great Parus major Great Dendrocopos major Great Wooc Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	chaff ared Dove amon Gull a Bunting koo ew erel chaff ared Marbler larest len Plover lafinch at Black-backed	mixed flock seen in January 2005 (IW) 6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	NL8 PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, BRC PL, BRC
Streptopelia decaocto Larus canus Comr Emberiza calandra Corn Cuculus canorus Numenius arquata Charadrius morinellus Prunella modularis Turdus pilaris Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Great Wooc Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Gomma	ared Dove Imon Gull Bunting Koo ew erel hock Ifare den Warbler Icrest Ien Plover Ifinch at Black-backed	6%, only seen in June 2009 (PL) 8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC, BBBW PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, BRC PL, BRC
Streptopelia decaocto Larus canus Comr Emberiza calandra Corn Cuculus canorus Numenius arquata Charadrius morinellus Prunella modularis Turdus pilaris Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Great Wooc Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Gomma	ared Dove Imon Gull Bunting Koo ew erel hock Ifare den Warbler Icrest Ien Plover Ifinch at Black-backed	8% (PL) 28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, ISGJ PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, BRC PL, BRC PL, IW, BRC, ISGJ
Larus canus Comr Emberiza calandra Corn Cuculus canorus Cucke Numenius arquata Curle Charadrius morinellus Dotte Prunella modularis Dunn Turdus pilaris Fieldf Sylvia borin Garde Regulus regulus Golde Pluvialis apricaria Golde Carduelis carduelis Goldf Larus marinus Great Gull Otis tarda Great Parus major Great Dendrocopos major Great Wooc Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	mon Gull Bunting soo ew erel nock Ifare den Warbler lcrest len Plover Ifinch at Black-backed	28% (PL) 6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, BRC PL, BRC PL, IW, BRC, ISGJ
Emberiza calandra Cuculus canorus Cucke Numenius arquata Charadrius morinellus Prunella modularis Turdus pilaris Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Tringa nebularia Crev Cucke Cucke Curle Curle Carduelis Carduelis Golde Golde Golde Golde Golde Great Gull Great Wooc Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	Bunting Koo Ew Erel Mock Ifare Iden Warbler Iden Plover Ifinch at Black-backed	6% (PL), most sightings in 2007 and 2009 (PL); 3 seen in 2005 (IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC, ISGJ, BBBW PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, BRC PL, W, BRC, ISGJ
Cuculus canorus Cucke Numenius arquata Curle Charadrius morinellus Dotte Prunella modularis Turdus pilaris Sylvia borin Regulus regulus Pluvialis apricaria Garduelis carduelis Garduelis carduelis Great Gull Otis tarda Parus major Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Tringa nebularia Grey Grey Grey Grey Grey Grey Grey Greet Garduelis chloris Greer Greet Grey Greet Grey Greet Grey Greet	coo ew erel nock Ifare den Warbler lorest len Plover Ifinch at Black-backed	(IW); flock of 28 seen in 2007 (IW) 2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, ISGJ PL PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, W, BRC, ISGJ
Numenius arquata Curle Charadrius morinellus Dotte Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Goldc Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Great Wooc Picus viridis Greer Carduelis chloris Greer Tringa nebularia Grey Grey Grey Grey Grey Greer Grey	ew erel nock Ifare den Warbler lcrest len Plover Ifinch at Black-backed	2%, heard in 2004, 2005 and 2008 (PL) 1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, W, BRC, ISGJ
Numenius arquata Curle Charadrius morinellus Dotte Prunella modularis Turdus pilaris Fieldf Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Goldc Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Great Wooc Picus viridis Greer Carduelis chloris Greer Tringa nebularia Grey Grey Grey Grey Grey Greer Grey	ew erel nock Ifare den Warbler lcrest len Plover Ifinch at Black-backed	1%, only seen in April 2009 (PL) The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, W, BRC, ISGJ
Charadrius morinellus Prunella modularis Turdus pilaris Sylvia borin Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Otis tarda Parus major Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Tringa nebularia Grey Doubte Doubte Doubte Doubte Garde Golde	erel mock Ifare den Warbler Icrest Ien Plover Iffinch It Black-backed	The "West Field" [N of section 7] was noted as a stopping-off place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 5% (PL) Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC, ISGJ, BBBW PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, W, BRC, ISGJ
Prunella modularis Turdus pilaris Fieldf Sylvia borin Garde Regulus regulus Pluvialis apricaria Carduelis carduelis Larus marinus Great Gull Otis tarda Parus major Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Grey Grey Grey Grey Grey Greer Grey	nock Ifare Ien Warbler Icrest Ien Plover Ifinch It Black-backed	place for this species in the third quarter of the 19th century, although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, IW, BRC, ISGJ
Turdus pilaris Fieldf Sylvia borin Garde Regulus regulus Golde Pluvialis apricaria Garduelis carduelis Larus marinus Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Tringa nebularia Fieldf Forder Golde Golde Great Gull Otis tarda Great Wood Ficus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	lfare len Warbler lcrest len Plover lfinch at Black-backed	although it is not clear when they were last seen there (NL5) 39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, IW, BRC, ISGJ
Turdus pilaris Fieldf Sylvia borin Garde Regulus regulus Golde Pluvialis apricaria Garduelis carduelis Larus marinus Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Tringa nebularia Fieldf Forder Golde Golde Great Gull Otis tarda Great Wood Ficus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	lfare len Warbler lcrest len Plover lfinch at Black-backed	39% (PL) 41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, IW, BRC, ISGJ
Turdus pilaris Fieldf Sylvia borin Garde Regulus regulus Golde Pluvialis apricaria Garduelis carduelis Larus marinus Otis tarda Parus major Dendrocopos major Picus viridis Greet Carduelis chloris Greet Tringa nebularia Fieldf Forder Golde Golde Great Gull Otis tarda Great Wood Ficus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	lfare len Warbler lcrest len Plover lfinch at Black-backed	41% (PL), A flock of approx. 300 seen in bushes in 2005 (IW); a total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, IW, BRC, ISGJ
Sylvia borin Garde Regulus regulus Golde Pluvialis apricaria Golde Carduelis carduelis Goldf Larus marinus Great Gull Otis tarda Great Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Grey Ardea cinerea Golde	len Warbler Icrest Ien Plover Ifinch It Black-backed	total of over 800 seen in flocks along its length in 2007 1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC, ISGJ, BBBW PL, BRC PL, BRC PL, IW, BRC, ISGJ
Regulus regulus Goldo Pluvialis apricaria Goldo Carduelis carduelis Goldf Larus marinus Great Gull Otis tarda Great Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Grey Ardea cinerea Goldo	crest len Plover lfinch at Black-backed	1%, only seen in June 1999 and July 2004 (PL) 24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC PL, BRC PL, IW, BRC, ISGJ
Regulus regulus Goldo Pluvialis apricaria Goldo Carduelis carduelis Goldf Larus marinus Great Gull Otis tarda Great Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Grey Ardea cinerea Goldo	crest len Plover lfinch at Black-backed	24% (PL) 5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC PL, BRC PL, IW, BRC, ISGJ
Pluvialis apricaria Golde Carduelis carduelis Goldf Larus marinus Great Gull Otis tarda Great Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Grey Ardea cinerea Grey	en Plover Ifinch at Black-backed	5% (PL) 51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, BRC PL, IW, BRC, ISGJ
Carduelis carduelis Carduelis carduelis Carduelis carduelis Great Gull Otis tarda Great Parus major Dendrocopos major Picus viridis Carduelis chloris Tringa nebularia Ardea cinerea Great Grea	lfinch at Black-backed	51% (PL), Part of a mixed flock seen in January 2005 (IW) 16% (PL)	PL, IW, BRC, ISGJ
Larus marinus Great Gull Otis tarda Great Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greey Ardea cinerea Grey	t Black-backed	16% (PL)	
Gull Otis tarda Great Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey			PL
Otis tarda Great Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	nt Bustard	Leather the advantage of the AOther and a second and the beautiful to the AOTher and the beautiful to the AOTHER A	
Parus major Great Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	it Bustard		
Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey		In the third quarter of the 19th century, one was noted to have been shot near the "West Field" [N of section 7], although it is not clear when this was (NL5)	
Dendrocopos major Great Wood Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	st Tit	50% (PL)	PL, BRC, ISGJ, BBBW
Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey	ater Spotted	7% (PL)	PL. BRC. ISGJ
Picus viridis Greer Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey		/ /o (i =)	1 2, 5110, 1000
Carduelis chloris Greer Tringa nebularia Greer Ardea cinerea Grey		27% (PL)	PL, BRC
Tringa nebularia Greer Ardea cinerea Grey		36% (PL), Part of a mixed flock seen in January 2005 (IW)	PL, IW, ISGJ, BBBW
Ardea cinerea Grey	enshank	5575 (* 2);	BRC
	Heron	4% (PL)	PL
Sidy		The general area is noted as a "paradise for partridges" in the third quarter of the 19th century (NL5); 4% (PL); 5 seen in 2005 (IW); 3 seen in 2006 (IW)	PL, IW, BRC, ISGJ
Larus argentatus Herrir	ing Gull	15% (PL)	PL
Falco subbuteo Hobb		1 seen in 2005 (IW)	IW, BRC
	se Martin	7% (PL)	PL, ISGJ
	se Sparrow	11% (PL)	PL, ISGJ
Corvus monedula Jacko		Many in beech plantation in section 1b in 1950s (NL5)	ISGJ
Garrulus glandarius Jay		4% (PL)	PL, ISGJ
Falco tinnunculus Kestro	rel	38% (PL)	PL, BRC, ISGJ
Vanellus vanellus Lapwi	ving	18% (PL); "plentiful" in 1950s (NL5)	PL, BRC, BBBW
	er Whitethroat		BRC, BBBW
Carduelis cannabina Linne	et	3% (PL)	PL, BRC, ISGJ, BBBW
	g-tailed Tit	77% (PL)	PL, BRC, ISGJ
Pica pica Magp	pie	66% (PL)	PL, ISGJ
Anas platyrhynchos Malla		1%, only seen in April 2004 (PL)	PL
	sh Harrier		BRC
	dow Pipit		BRC, ISGJ
Turdus viscivorus Mistle	e Thrush		BRC
Phasianus colchicus Pheas		67% (PL)	PL, ISGJ
	Wagtail	4% (PL)	PL
Coturnix Quail		1 heard at the Balsham end in 2009 (IW)	IW, BRC
	backed Shrike	2 pairs seen in 1941 (NL9), "occasional visitors" in 1950s (NL5)	
Alectoris rufa Red-l	legged Partridge	82% (PL)	PL, BRC, ISGJ
Turdus iliacus Redw		33% (PL), A flock of about 50 seen with Fieldfare in 2005 (IW)	PL, IW, BRC
		100 /0 (1 E); / (1100 k 0 1 about 00 000 h with 1 10 and a 10 h 2000 h w	PL
		1%, only seen in June 2005 and March 2009 (PL)	· -
Erithacus rubecula Robin	wing		BRC

Corvus frugilegus	Rook	58% (PL)	PL, ISGJ
Asio flammeus	Short-eared Owl	3% (PL)	PL
Carduelis spinus	Siskin	1 seen in 2002 (NL7); 1 seen in 2003 (NL8)	NL7, NL8
Alauda arvensis	Skylark	66% (PL), "plentiful" in 1950s (NL5)	PL, BRC, ISGJ, BBBW
Turdus philomelos	Song Thrush	6% (PL)	PL, BRC, ISGJ, BBBW
Accipiter nisus	Sparrowhawk	40% (PL), 1 seen in 2005 (IW)	PL, IW, BRC, ISGJ
Muscicapa striata	Spotted Flycatcher	1%, only seen in July 2002 (PL)	PL, BRC
Sturnus vulgaris	Starling	11% (PL), A flock of 150 seen in 2005 (IW)	PL, IW, BRC, ISGJ
Burhinus oedicnemus	Stone Curlew	Seen feigning injury on land belonging to Grange Farm, Little	
		Abington, in 1948 (NL9); one seen in fields at back of Wandlebury	
		in 1950s (NL5)	
Hirundo rustica	Swallow	34% (PL), 1 seen in April 2007 (IW)	PL, IW, BRC, BBBW
Apus apus	Swift	13% (PL)	PL, ISGJ
Strix aluco	Tawny Owl	2% (PL)	PL
Streptopelia turtur	Turtle Dove	9% (PL)	PL, BRC, BBBW
Oenanthe oenanthe	Wheatear	3% (PL)	PL
Sylvia communis	Whitethroat	23% (PL)	PL, BRC, BBBW
Phylloscopus trochilus	Willow Warbler	10% (PL)	PL, BRC, BBBW
Columba palumbus	Wood Pigeon	88% (PL)	PL, BRC, ISGJ
Troglodytes troglodytes	Wren	48% (PL)	PL, BRC, ISGJ, BBBW
Jynx torquilla	Wryneck	"Occasional visitors" in 1950s (NL5)	
Motacilla flava	Yellow Wagtail	2 flew over in 2005 (IW)	IW, BRC
Emberiza citrinella	Yellowhammer	69% (PL)	PL, BRC, ISGJ, BBBW